

Increased Response of 5xFAD Mouse Primary Adult Microglia to Different Pro-Inflammatory Stimuli

Roland Rabl, Irene Schilcher, Tina Loeffler, Birgit Hutter-Paier, Manuela Prokesch
QPS Austria, Parkring 12, 8074 Grambach, Austria

BACKGROUND

The importance of microglia in neurodegenerative diseases is well-known and these cells are therefore frequently used as target for new pharmacological interventions. To study this cell type, isolation of early postnatal microglia from mice is a great tool but does not properly reflect conditions in aged or diseased individuals. Isolation of viable microglia from adult mouse brains of specific disease models via Magnetic Cell Sorting (MACS) opens new opportunities to assess the efficacy of microglia-targeting treatments *in vitro*. Here we investigated the *in vitro* response of isolated adult microglia from 5xFAD mice to various stimuli in comparison to age-matched non-transgenic (nTg) microglia.

MATERIAL & METHODS


Microglia were isolated from 9-months old 5xFAD and nTg mice. Cultivated cells were stimulated with LPS or Aβ1-42 in presence or absence of the anti-inflammatory agent dexamethasone. Release of the cytokines TNF-α, IL-6, IL-1β and KC/GRO, as well as Aβ1-42 phagocytosis and Aβ1-42 levels in the supernatant were measured.

RESULTS


While a strong enrichment of Itgam (CD11b) positive cells in the microglial fraction compared to the flow through was detected, GFAP expression was extremely low or absent.

Figure 1: Purity of isolated microglia. RT-PCR amplification graphs of Itgam (A) and GFAP (B) SYBR green signal in microglia and flow through fractions prepared from 3 separate 9.5 months old 5xFAD mouse brains. The graphs show relative fluorescence units (RFU) of the SYBR green signal per cycle.

A Itgam (CD11b)



B GFAP



The response to pro-inflammatory stimuli was tremendously stronger in 5xFAD microglia, revealing a 10-fold higher level of secreted cytokines compared to nTg microglia that were stimulated with the same stressor. Treatment of LPS-stimulated 5xFAD microglia with the anti-inflammatory agent dexamethasone was able to significantly attenuate the cytokine release compared to LPS-stimulated control cells.

LPS-Induced Inflammation




Figure 2: Quantification of TNF-α, IL-6 and IL-1β levels in the supernatant of LPS-stimulated microglia.

Isolated microglia of 5xFAD mice and nTg littermates were treated with Vehicle (VC), 50 ng/mL LPS or LPS plus dexamethasone (Dexa) and TNF-α (A), IL-6 (B) and IL-1β (C) secretion was measured. Data are presented as bar graphs with mean + SEM (n=4-6 per group). Two-Way ANOVA followed by Bonferroni's Multiple Comparison *post hoc* test compared to LPS control; *p<0.01; ***p<0.001 or nTg versus 5xFAD ###p<0.001.

Aβ1-42-Induced Inflammation and Phagocytosis




Figure 3: Quantification of TNF-α, IL-6 and IL-1β levels in the supernatant as well as phagocytosis of Aβ1-42-treated microglia.

Isolated microglia of 5xFAD mice and nTg littermates were treated with Vehicle (VC) or 10 μM Aβ1-42 and TNF-α (A), IL-6 (B) and IL-1β (C) secretion as well as phagocytosis (D,E) and Aβ1-42 levels in the supernatant (F) were measured. Data are presented as bar graphs with mean + SEM (n=5 per group). (A-C) Two-Way ANOVA followed by Bonferroni's Multiple Comparison *post hoc* test; (E-F) Two-tailed unpaired *t*-test; ***p<0.001 or nTg versus 5xFAD ###p<0.001.

SUMMARY and CONCLUSION

Generation of a pure microglial fraction from adult brains of transgenic mice opens a variety of new opportunities to assess the efficacy of treatments in diseased microglia.

Meet QPS at our Booth

For more information about the models please visit:

www.qpsneuro.com or send us an e-mail: office-austria@qps.com